



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

to be related to the character of the natural habitat of the species in question. Thus *Salix* sp. (probably *nigra*) stands at one end of the series and shows no injurious effect even when the oxygen of the atmosphere is entirely replaced by either nitrogen or carbon dioxide. At the opposite end of the series stands *Opuntia versicolor*, growth of roots ceasing with an atmosphere containing 50 per cent carbon dioxide, while *Coleus Blumei* is comparable to it, showing injury and ultimate death with the addition of 25 per cent nitrogen to the soil atmosphere. Of the other species tested *Heliotropium peruvianum* was closely comparable to *Opuntia*, while *Nerium oleander* and *Prosopis velutina* prove nearly as resistant as *Salix*. The results seem to indicate that plants growing naturally in well drained soil are much more sensitive to the composition of the soil atmosphere than those from swamps and poorly drained habitats.—GEO. D. FULLER.

Embryo of *Aucuba*.—PALM and RUTGERS³¹ have settled the question of apogamy in *Aucuba japonica*, which has been under suspicion for 40 years. They bagged 300 pistillate flowers and not a single fruit formed, while 600 isolated pistillate flowers produced normal fruit after artificial pollination. It is thought that EICHLER's original suggestion of apogamy probably came from the fruiting of an isolated pistillate plant which had developed staminate flowers, since the authors have repeatedly found staminate flowers on pistillate plants. Staminate plants have also been observed to produce pistillate flowers.

The flowers open about the time of megaspore formation, and the embryo sac reaches the fertilization stage about 4 weeks later. The solitary megaspore mother cell becomes deeply placed by the extensive development of parietal tissue. The behavior of the 4 megaspores is usually quite normal, but in one case the 2 megaspores nearest the chalaza were found in division. The development of the gametophyte is normal, but stages in endosperm formation were not obtained. The chromosome numbers were determined to be 18 and 36.—J. M. C.

Disease resistance.—JONES³² has published a summary of his results in securing a race of cabbage resistant to the "yellows." Some of the fundamental questions involved in resistance were considered. The difference between susceptible and resistant plants was found not to be due to any superficial obstacle, but to the different relations of the interior cells of the host and parasite. "The resistant tissues have the ability to restrain the development of the parasite to a greater degree than do the susceptible and so give time for protective cork formation." It was shown also that resistance is clearly inheritable, not as a single character, but as a complex of a number of heritable

³¹ PALM, BJ., and RUTGERS, A. A. L., The embryology of *Aucuba japonica*. Rec. Trav. Bot. Néerland. 14:119-126. figs. 12. 1917.

³² JONES, L. R., Disease resistance in cabbage. Proc. Nat. Acad. Sci. 4:42-46. 1918.